

Cosmological Particle Creation in States of Low Energy

Andreas Degner and Rainer Verch

Institut für Theoretische Physik
Universität Leipzig
Postfach 100 920
D-04009 Leipzig, Germany

Abstract

The recently proposed states of low energy provide a well-motivated class of reference states for the quantized linear scalar field on cosmological Friedmann-Robertson-Walker spacetimes. The low energy property of a state is localized close to some value of the cosmological time coordinate. We present calculations of the relative cosmological particle production between a state of low energy at early time and another such state at later time. In an exponentially expanding Universe, we find that the particle production shows oscillations in the spatial frequency modes. The basis of the method for calculating the relative particle production is completely rigorous. Approximations are only used at the level of numerical calculation.

1 Introduction

The phenomenon of particle production on non-stationary cosmological spacetimes is one of the early findings in the theory of quantized fields on curved spacetimes [17]. At its very basics, it can be reduced to an analogy between the equation of motion for the mode functions of a quantum field in a time-dependent (spatially homogeneous) spacetime metric and the equation of motion for a harmonic oscillator with time-dependent frequency [17, 8]. It is, however, not easy to make the analogy completely rigorous, for the mode decomposition on which the notion of particles is based is itself time-dependent.

Quite generally, in a time-dependent background, any concept of particle will itself depend on time, and possibly on other data, such as scale parameters. In more precise terms, for a quantum field on a spacetime with time dependent metric, a concept of particle is tied to the selection of a reference state, which then depends on such data as a hypersurface of constant time, and other, scale-setting parameters. In the present paper, we will be concerned with the case of a linear, minimally coupled scalar field on Friedmann-Robertson-Walker (FRW) spacetimes. For this situation, particle creation with respect to some classes of reference states has been investigated. Notably, Parker introduced so-called “adiabatic vacuum states”, designed to minimize particle creation, as reference states [17, 8]. The precise mathematical definition of this class of states is intricate and was first achieved in an investigation by Lüders and Roberts [15]. In further work by Junker and Schrohe [13], it was demonstrated that adiabatic vacua are locally unitarily equivalent to the class of Hadamard states. Since many results have, over the past decades, confirmed the view originally put forward by Wald [23, 25] that the class of physical states for linear quantum fields on curved spacetimes should contain all (quasifree) Hadamard states as building blocks, this shows that adiabatic vacua qualify as physical states as

well. However, the physical interpretation of adiabatic vacua remains less clear; furthermore, their definition involves, in the general case, asymptotic series expansions making them inconvenient to handle and posing difficulties when trying to obtain numerical results on particle creation [27].

Quite recently, Olbermann [16] has introduced a new class of reference states for the linear scalar field on FRW spacetimes, called *states of low energy* (SLEs). These states are spatially homogeneous (with respect to the spatial isometry group of FRW spacetimes) and such that they minimize the time-integral of the energy-density for a given weighting function. For the moment, let us describe this in more detail as follows (the exact definition will be given in Sec. 3). Suppose that t is the cosmological time coordinate in an FRW spacetime, and let $f(t)$ denote a non-negative, compactly supported, smooth function of cosmological time. For a homogeneous state of the quantized linear scalar field on FRW spacetime, denoted by its expectation functional $\langle \cdot \rangle_\omega$ where ω is a label for the state, we write $\langle : \varrho(t) : \rangle_\omega$ for the (renormalized) expected energy density in that state. As the state is homogeneous, this quantity is only dependent on time t . The time-integral of the expected renormalized energy density, weighted with respect to f , is

$$\varrho_\omega[f] = \int \langle : \varrho : (t) \rangle_\omega f(t) dt.$$

Then a state $\langle \cdot \rangle_{\omega_{\text{SLE}}}$ is called an *SLE (state of low energy) with respect to the weighting function f* if the state minimizes the expression $\varrho_\omega[f]$ among all homogeneous states $\langle \cdot \rangle_\omega$. Note that the SLE property depends on the choice of weighting function f and that it is therefore a concept that is local in time. Olbermann has shown that SLEs are Hadamard states, and that adiabatic vacua are approximations of SLEs.

If a state $\langle \cdot \rangle_\omega$ were an SLE for the case $f = \delta_{t_0}$, the delta-distribution concentrated at cosmological time t , it would correspond to a state minimizing the (homogeneous) energy density at time t_0 . This is reminiscent of the concept of an instantaneous vacuum state at time t_0 , and the interpretation of SLEs is that of approximate (or, by weighting with respect to f , mollified) instantaneous vacua at time t_0 . Instantaneous vacua, however, aren't Hadamard states, and they are not even (locally) unitarily equivalent to Hadamard states [9, 12], thus disqualifying them from the class of physical states. The weighting, or mollifying, of the energy density by integrating in time against smooth weighting functions f in defining SLEs is therefore needed in order to ensure that SLEs are physical states.

Nevertheless, the interpretation as homogeneous, energy minimizing states close to an instant of cosmological time makes SLEs natural candidates for reference states with respect to which particle concepts can be defined and particle creation in FRW spacetimes can be calculated. This is the topic of the present work. More specifically, we consider two smooth weighting functions, f_1 and f_2 , peaked at cosmological time parameters t_1 and t_2 , respectively. We will envisage the situation that t_1 corresponds to an early cosmological time and $t_2 > t_1$ a later cosmological time. Denoting by $\langle \cdot \rangle_{\omega(1)}$ and $\langle \cdot \rangle_{\omega(2)}$ the corresponding SLEs, we will determine the particle content per (fixed) frequency mode k (arising from the spatial symmetry) of the states with respect to each other; this quantity is given by the modulus squared $|\beta_k|^2$ of the Bogoliubov coefficients, see Sec. 4. The numerical calculation, however, turns out to be

involved, and therefore we limit ourselves in this work to a special case which simplifies the calculations significantly. However, we think that the principal qualitative aspects of the results of our calculations won't depend very critically on these choices. Our specialized assumptions are: (i) A closed (spatially compact) FRW spacetime, (ii) an exponentially increasing scale-factor $a(t) = e^{Ht}$ ($H > 0$), (iii) a very small mass parameter $m > 0$, fine-tuned with respect to the Hubble-parameter H , in the linear field equation of the quantized scalar field. Moreover, we approximate the weighting functions f_1 and f_2 by Gaussians peaked at t_1 and t_2 with small characteristic widths ϵ and δ , respectively. Then $|\beta_k|^2$ will be calculated numerically and represented graphically for certain choices of t_1, t_2, ϵ and δ .

The most interesting result is that $|\beta_k|^2$ shows oscillations in k when $t_2 - t_1$ is different from 0 and this effect increases with growing $t_2 - t_1$. A somewhat related observation to the effect that the particle number per mode can decrease in time appears in [11]; however, the reference states used in [11] are instantaneous vacua.

It is worth pointing out that our approach is completely rigorous, with approximations entering only at the level of numerical calculation. While the simplifications (i), (ii) and (iii) made in order to facilitate the numerical calculations are not physically realistic, it is, as mentioned, to be expected that the basic findings of our calculations, especially the said oscillatory behaviour of $|\beta_k|^2$, will qualitatively also occur in physically more realistic situations. This could have some implications with regard to observations in cosmology.

This article is organized as follows. The quantization of the linear (minimally coupled) scalar field on FRW spacetimes will be summarized in Sec. 2, together with the definition of the renormalized stress-energy tensor. States of low energy will be discussed in Sec. 3. In Sec. 4 we consider Bogoliubov transformations between two SLEs and the associated notion of particle creation. The numerical calculations will be presented in Sec. 5. Summary and Outlook are given in the final Sec. 6.

2 Quantum field on FRW spacetimes and renormalized stress-energy tensor

In this section, we summarize the quantization of the linear scalar Klein-Gordon field on FRW spacetimes, and the definition of the renormalized stress-energy tensor. This serves mainly to make the present text as self-contained as possible. General references for the material in this section are [23, 24, 25, 8].

A spacetime (M, g) , where M denotes the spacetime manifold and g the Lorentzian metric, is of FRW type if it is of the form

1. $M = \mathbb{R} \times \Sigma$
2. $ds_g^2 = dt^2 - a(t)^2 h_{ij} d\mathbf{x}^i d\mathbf{x}^j$
3. (Σ, h) is a 3-dimensional manifold with Riemannian metric of constant curvature which is either equal to 0 or normalized to ± 1 . (We write usually x for elements in M , identified also as $(t, \mathbf{x})$ with $t \in \mathbb{R}$, $\mathbf{x} \in \Sigma$.)
4. $a : \mathbb{R} \rightarrow \mathbb{R}$ is a C^∞ function taking strictly positive values, called the scale factor.

The classical linear, minimally coupled scalar field (Klein-Gordon field) φ on (M, g) fulfills the field equation

$$(\square_g + m^2)\varphi = 0$$

with smooth $\varphi : M \rightarrow \mathbb{R}$. The constant parameter $m \geq 0$ is called the mass term of the field equation and $\square_g$ is the d'Alembertian of (M, g) which in the present case takes the form

$$\square_g + m^2 = \frac{\partial^2}{\partial t^2} + 3H(t)\frac{\partial}{\partial t} - \frac{1}{a(t)^2}\Delta_h + m^2. \quad (1)$$

Here, $H(t) = (\frac{d}{dt}a(t))/a(t)$ is the Hubble function, and Δ_h denotes the Laplace operator on (Σ, h) . An FRW spacetime (M, g) is a globally hyperbolic spacetime and therefore there exists a unique pair, $E^\pm$, of advanced/retarded Green's functions for the Klein-Gordon operator $\square_g + m^2$, for any fixed $m \geq 0$ [5, 1]. They are characterized as continuous linear maps $E^\pm : C_0^\infty(M, \mathbb{R}) \rightarrow C^\infty(M, \mathbb{R})$ with $E^\pm(\square_g + m^2)F = F = (\square_g + m^2)E^\pm F$ for all $F \in C_0^\infty(M, \mathbb{R})$, and with $\text{supp}(E^\pm F) \subset J^\pm(\text{supp } F)$ for all $F \in C_0^\infty(M, \mathbb{R})$, where $J^\pm(G)$ denotes the future(+)/past(-) causal set of a subset $G \subset M$. The difference $E = E^+ - E^-$ is often called causal propagator of the linear scalar field on (M, g) . We write

$$\mathcal{E}(F_1, F_2) = \int_M F_1(x)(EF_2)(x) d\mu_g(x) \quad (F_1, F_2 \in C_0^\infty(M, \mathbb{R}))$$

for the associated bilinear form, where $d\mu_g(x) = \sqrt{|\det(g_{\mu\nu})|}d^4x$ is the metric-induced volume form on M . One can show that $\mathcal{E}$ is antisymmetric, i.e. $\mathcal{E}(F_1, F_2) = -\mathcal{E}(F_2, F_1)$.

The linear scalar field on (M, g) can then be quantized as follows. One defines an abstract $*$ -algebra $\mathcal{F}(M, g)$ with algebraic unit element $\mathbf{1}$ as being generated by $\mathbf{1}$ and a family of elements $\phi(F)$, $F \in C_0^\infty(M, \mathbb{R})$, with the relations

$$\phi(F_1)\phi(F_2) - \phi(F_2)\phi(F_1) = \frac{i}{2}\mathcal{E}(F_1, F_2),$$

$$\phi(F)^* = \phi(F),$$

$$\phi((\square_g + m^2)F) = 0$$

for all $F, F_1, F_2 \in C_0^\infty(M, \mathbb{R})$.

One can show that such an algebra exists (nontrivially) and that it is unique up to (natural) isomorphisms (cf. e.g. [2] and lit. cited there). The generating elements $\phi(F)$ can be viewed as “abstract field operators” for the quantized linear scalar field on (M, g) . The $\phi(F)$ can be turned into operators acting in a Hilbert space upon considering Hilbert space representations of $\mathcal{F}(M, g)$. Here, we are interested in Hilbert space representations induced by states on $\mathcal{F}(M, g)$. Recall that a state on $\mathcal{F}(M, g)$ is a linear functional $\omega : \mathcal{F}(M, g) \rightarrow \mathbb{C}$ which is positive, i.e. it fulfills $\omega(X^*X) \geq 0$ for all $X \in \mathcal{F}(M, g)$. (Note that X is in $\mathcal{F}(M, g)$ if it is a polynomial built out of $\mathbf{1}$ and finitely many $\phi(F_1), \dots, \phi(F_N)$.) Moreover, it is required that ω be continuous, that is, for each $n \in \mathbb{N}$ the map $F_1 \otimes \dots \otimes F_n \mapsto \omega(\phi(F_1) \dots \phi(F_n))$ extends to a distribution in $(C_0^\infty(M^n, \mathbb{R}))'$. It is also common to denote a state by its expectation value functional, $\langle X \rangle_\omega = \omega(X)$, and we shall often adopt this notation.

There are states ω entirely determined by their 2-point function

$$\mathcal{W}_2^\omega(F_1, F_2) = \omega(\phi(F_1)\phi(F_2)) = \langle \phi(F_1)\phi(F_2) \rangle_\omega \quad (F_1, F_2 \in C_0^\infty(M, \mathbb{R}))$$

by requiring the relations $\langle \phi(F_1) \cdots \phi(F_{2n+1}) \rangle_\omega = 0$ and

$$\left. \frac{d^{2n}}{d\lambda^{2n}} \right|_{\lambda=0} \langle e^{i\lambda\phi(F)} \rangle_\omega = \left. \frac{d^{2n}}{d\lambda^{2n}} \right|_{\lambda=0} e^{-\lambda W_2^\omega(F,F)/2} \quad (F \in C_0^\infty(M, \mathbb{R}))$$

for all $n \in \mathbb{N}$; the left hand side of this equation is to be read as the expectation value of the polynomial in $\phi(F)$ that results from formally differentiating $e^{i\lambda\phi(F)}$ and setting λ equal to 0.

A state ω on $\mathcal{F}(M, g)$ induces a $*$ -representation π_ω of $\mathcal{F}(M, g)$ on a dense domain $\mathcal{D}_\omega$ in a Hilbert space $\mathcal{H}_\omega$ together with a canonical unit vector $\Omega_\omega \in \mathcal{D}_\omega$. The collection of $(\mathcal{H}_\omega, \mathcal{D}_\omega, \pi_\omega, \Omega_\omega)$ is called GNS-representation, or Wightman-representation of ω (cf. [21]) and is characterized by the properties that $\langle X \rangle_\omega = \langle \Omega_\omega, \pi_\omega(X) \Omega_\omega \rangle$ for all $X \in \mathcal{F}(M, g)$, $\pi_\omega(X^*)\chi = \pi_\omega(X)^*\chi$ for all $\chi \in \mathcal{D}_\omega$, and $\mathcal{D}_\omega = \{\pi_\omega(X)\Omega_\omega : X \in \mathcal{F}(M, g)\}$. Then the represented abstract field operators $\phi(f)$ become unbounded operators

$$\Phi_\omega(f) = \pi_\omega(\phi(f)) \quad (2)$$

defined on the domain $\mathcal{D}_\omega$ in the representation Hilbert space $\mathcal{H}_\omega$.

For a quasifree state ω , there is a one-particle Hilbert space $\mathcal{H}_\omega^{(1)}$ together with a real-linear map $K_\omega : C_0^\infty(M, \mathbb{R}) \rightarrow \mathcal{H}_\omega^{(1)}$ so that $\langle K_\omega(F_1), K_\omega(F_2) \rangle = W_2^\omega(F_1, F_2)$. One can then show that $\mathcal{H}_\omega = F_+(\mathcal{H}_\omega^{(1)})$, the bosonic Fock-space over the one-particle space $\mathcal{H}_\omega^{(1)}$, and that

$$\pi_\omega(\phi(F)) = a(K_\omega(F)) + a^+(K_\omega(F)) \quad (3)$$

where $a(\cdot)$ and $a^*(\cdot)$ are the usual annihilation and creation operators on $F_+(\mathcal{H}_\omega^{(1)})$. Furthermore, $\Omega_\omega = (1, 0, 0, \dots)$ is the Fock-vacuum-vector and thus it is convenient to write $|0\rangle_\omega = \Omega_\omega$ for the GNS-vector of a quasifree states ω . Moreover, a quasifree state ω is *pure* if the range of K_ω is dense in $\mathcal{H}_\omega^{(1)}$. This is equivalent to saying that ω cannot be written as convex combination of several different states.

On each FRW spacetime $(M = \mathbb{R} \times \Sigma, g)$ there acts the spatial symmetry group G_Σ . It acts only in the Σ -part of M , so that for each $\gamma \in G_\Sigma$ one has $\gamma(t, \mathbf{x}) = (t, \gamma_\Sigma(\mathbf{x}))$ for all $t \in \mathbb{R}$, $\mathbf{x} \in \Sigma$ with respect to the $\mathbb{R} \times \Sigma$ -splitting of M , with an isometry γ_Σ of (Σ, h) .¹ By setting $\alpha_\gamma(\phi(F)) = \phi(F \circ \gamma^{-1})$ for $F \in C_0^\infty(M, \mathbb{R})$ and $\gamma \in G_\Sigma$, one can define the automorphisms $\alpha_\gamma : \mathcal{F}(M, g) \rightarrow \mathcal{F}(M, g)$ induced by γ on the algebra of abstract field operators. Then a state ω on $\mathcal{F}(M, g)$ is called *homogeneous* if it is invariant under the action of α_γ , i.e. if

$$\langle \alpha_\gamma(X) \rangle_\omega = \langle X \rangle_\omega \quad (\gamma \in G_\Sigma, X \in \mathcal{F}(M, g)).$$

If ω is a quasifree state, then it is homogeneous exactly if

$$W_2^\omega(F_1 \circ \gamma^{-1}, F_2 \circ \gamma^{-1}) = W_2^\omega(F_1, F_2)$$

holds for all $\gamma \in G_\Sigma$ and $F_1, F_2 \in C_0^\infty(M, \mathbb{R})$.

The most important class of states for quantized linear fields on curved spacetimes, particularly in the context of defining expectation values of the

¹Strictly speaking, $G_\Sigma = \{\text{id}_\mathbb{R} \times \gamma_\Sigma : \gamma_\Sigma \in \text{Iso}^+(\Sigma, h)\}$ where $\text{Iso}^+(\Sigma, h)$ is the Lie-group of orientation-preserving isometries of (Σ, h) .

stress-energy tensor, are quasifree Hadamard states. These are quasifree states whose 2-point functions are of so-called Hadamard form. We shall not go into full details of the definition of Hadamard form (see [14, 25, 7]) for further discussion) but mainly describe its basic features entering into the definition of the renormalized expected stress-energy tensor. Basically, $\mathcal{W}_2^\omega$ is of Hadamard form if

$$\mathcal{W}_2^\omega(F_1, F_2) = \mathcal{G}_{\text{sing}}(F_1, F_2) + \int \int F_1(x) R_\omega(x, x') F_2(x') d\mu_g(x) d\mu_g(x')$$

where $\mathcal{G}_{\text{sing}}$ is a singular contribution (a distribution in $(C_0^\infty(M \times M))'$) which depends in a certain, local way on the spacetime metric g and the mass parameter m in the Klein-Gordon equation, but is independent of ω (i.e. the singular part is the same for all Hadamard states ω). There remains a smooth contribution, expressed by $R_\omega \in C^\infty(M \times M, \mathbb{C})$, which contains the dependence on the states ω . The circumstance that the singular part $\mathcal{G}_{\text{sing}}$ is the same for all Hadamard states ω is instrumental for the definition of the expectation value $\langle : \mathbf{T}_{\mu\nu} : \rangle_\omega$ of the renormalized stress-energy tensor in the state ω .

We shall very briefly elaborate on the “symmetric Hadamard parametrix” (SHP) renormalization of the expectation value of the stress-energy tensor which was employed in [19], see also [7]. Define the *symmetric Hadamard parametrix*

$$\tilde{\mathcal{G}}(F_1, F_2) = \frac{1}{2}(\mathcal{G}_{\text{sing}}(F_1, F_2) + \mathcal{G}_{\text{sing}}(F_2, F_1)) + \frac{i}{2}\mathcal{E}(F_1, F_2)$$

and, correspondingly, set

$$\mathcal{R}_\omega^{\text{SHP}}(F_1, F_2) = \mathcal{W}_2^\omega(F_1, F_2) - \tilde{\mathcal{G}}(F_1, F_2)$$

for $F_1, F_2 \in C_0^\infty(M, \mathbb{R})$. Then the distribution $\mathcal{R}_\omega^{\text{SHP}}$ is given by a smooth integral kernel R_ω^{SHP} ,

$$\mathcal{R}_\omega^{\text{SHP}}(F_1, F_2) = \int \int F_1(y) R_\omega^{\text{SHP}}(x, x') F_2(x') d\mu_g(x) d\mu_g(x').$$

Now let $x \in M$ be given and suppose that $x' \in M$ lies in a convex normal neighbourhood of x , and denote by $Y^{\nu'}_\nu(x'; x)$ the operation of parallelly transporting a covector $\xi'_{\nu'}$ at x' to a covector ξ_ν at x . Note that $Y^{\nu'}_\nu(x'; x)$ depends smoothly on x and x' . With this convention, one can define the SHP-renormalized expectation value of the stress-energy tensor in the state ω as

$$\begin{aligned} \langle \mathbf{T}_{\mu\nu}^{\text{SHP}}(x) \rangle_\omega &= \lim_{x' \rightarrow x} (\nabla_\mu \nabla_{\nu'} R_\omega^{\text{SHP}}(x, x')) Y^{\nu'}_\nu(x'; x) \\ &\quad + \frac{1}{2} g^{\mu\nu}(x) (\nabla_\mu \nabla_{\nu'} R_\omega^{\text{SHP}}(x, x')) Y^{\nu'}_\nu(x'; x) \\ &\quad + \frac{1}{2} R_\omega^{\text{SHP}}(x, x'). \end{aligned}$$

Note that on the right hand side, ∇_μ operates with respect to x and $\nabla_{\nu'}$ operates with respect to x' . The resulting $\langle \mathbf{T}_{\mu\nu}^{\text{SHP}}(x) \rangle_\omega$ is a smooth, symmetric co-tensor field of $x \in M$. This renormalization procedure has the advantage of being completely independent of any “reference” state; in fact, it renders a local, generally covariant quantity in the sense of [25, 2, 10]. However, in general it

will have the defect of not being divergence-free. As pointed out in [23, 25], this defect can be repaired as follows: One can show that there is a smooth function Q on M , determined entirely by the local geometry of (M, g) , such that

$$\nabla^\mu \langle \mathbf{T}_{\mu\nu}^{\text{SHP}}(x) \rangle_\omega = \nabla_\nu Q(x) \quad (x \in M)$$

holds for all quasifree Hadamard states ω – while Q is state-independent. Hence, if $Q(x)g_{\mu\nu}(x)$ is subtracted from $\langle \mathbf{T}_{\mu\nu}^{\text{SHP}}(x) \rangle_\omega$, the resulting quantity has vanishing divergence. In fact, this resulting quantity is a local, generally covariant and divergence-free definition of the expectation value of the stress-energy tensor of a quantized field which thus complies with the requirements delineated by Wald for stress-energy observables of quantum fields [23, 25]. Then there remains a renormalization ambiguity for this quantity to the effect that one may add other symmetric co-tensor fields $C_{\mu\nu}$ having vanishing divergence and determined by the local geometry of (M, g) . Following [7], we adopt the point of view that the specification of $C_{\mu\nu}$ is a further datum (akin to the mass parameter m) of the quantum field ϕ . Using an argument of [25], the freedom of choosing $C_{\mu\nu}$ can be further narrowed down, so that

$$C_{\mu\nu} = C_{\mu\nu}[A, B, \Gamma, D] = Ag_{\mu\nu} + BG_{\mu\nu} + \Gamma \frac{\delta}{\delta g^{\mu\nu}} S_1(g) + D \frac{\delta}{\delta g^{\mu\nu}} S_2(g),$$

where $G_{\mu\nu}$ is the Einstein tensor, $S_1(g) = \int_M R^2 d\mu_g$, $S_2(g) = \int_M R_{\mu\nu} R^{\mu\nu} d\mu_g$ with R and $R_{\mu\nu}$ denoting scalar curvature and Ricci tensor, while $\delta/\delta g^{\mu\nu}$ denotes functional differentiation; the real constants A, B, Γ and D parametrize the remaining renormalization ambiguity. In conclusion, the renormalized expected stress-energy tensor in a quasifree Hadamard state ω takes the form

$$\langle : \mathbf{T}_{\mu\nu} : (x) \rangle_\omega = \langle \mathbf{T}_{\mu\nu}^{\text{SHP}}(x) \rangle_\omega - Q(x)g_{\mu\nu}(x) + C_{\mu\nu}[A, B, \Gamma, D](x) \quad (x \in M).$$

Considering the case of an FRW spacetime with cosmological time coordinate t , the renormalized expected energy density in a quasifree Hadamard state ω with respect to cosmological time is

$$\langle : \rho : (t, \mathbf{x}) \rangle_\omega = \langle : \mathbf{T}_{\mu\nu} : (t, \mathbf{x}) \rangle_\omega \left(\frac{\partial}{\partial t} \right)^\mu \left(\frac{\partial}{\partial t} \right)^\nu \quad ((t, \mathbf{x}) \in \mathbb{R} \times \Sigma).$$

It is worth making a few remarks here.

(i) The notation using double dots is to be understood as signifying that $\langle : \mathbf{T}_{\mu\nu} : (x) \rangle_\omega$ is a renormalized quantity. It is not to be confused with the more common usage of indicating “normal ordering”, which refers to a reference state. The same applies to our notation of $\langle : \rho : (x) \rangle_\omega$ for the renormalized expected energy density.

(ii) The above indicated procedure of defining the renormalized stress-energy expectation value of a linear quantized field in Hadamard states ω applies not only to the case of FRW spacetimes but to general, globally hyperbolic spacetimes.

(iii) It follows from the appearance of renormalization ambiguities in the definition of $\langle : \mathbf{T}_{\mu\nu} : (x) \rangle_\omega$ (parametrized by renormalization constants A, B, Γ, D) that there is no intrinsic (without further considerations) prediction of the total absolute value of the (expected) local energy density in quantum field theory on generic spacetimes prior to fixing the renormalization ambiguities. The way

in which the fixing is done can have significant implications in cosmological scenarios (see, e.g., [3]). In the context of discussions of the role played by the “vacuum energy” as contribution to the cosmological constant it is occasionally claimed that quantum field theory were predicting a value of the vacuum energy which misses the observed value by 120 orders of magnitude. It is worth pointing out that quantum field theory doesn’t make any such prediction.

3 States of low energy (SLEs)

Since our investigation later in this article refers to the case of spatially compact FRW spacetime, we will from now on restrict our discussion to that case, mostly to simplify notation. However, states of low energy have been introduced in [16] for all types of FRW spacetimes.

The spatially compact FRW spacetime has $\Sigma = S^3$ with the Riemannian metric h on Σ derived from the embedding

$$\Sigma = \{(y^1, y^2, y^3, y^4) \in \mathbb{R}^4 : \sum_{j=1}^4 (y^j)^2 = 1\}$$

into $\mathbb{R}^4$ equipped with the Euclidean metric. This Riemannian manifold carries the metric-induced measure $d\mu_h$ which in turn gives rise to the Hilbert space $L^2(\Sigma, d\mu_h)$. The Laplacian Δ_h is essentially selfadjoint on $C^\infty(\Sigma, \mathbb{C})$ and there is an orthonormal basis $\{Y_{\mathbf{k}}\}$, $\mathbf{k} = (k, l, m)$, $k \in \mathbb{N}_0$, $l = 0, 1, \dots, k$, $m = -l, -l+1, \dots, l$, of C^∞ functions on Σ which are eigenvectors of $-\Delta_h$,

$$-\Delta_h Y_{\mathbf{k}} = \kappa(\mathbf{k})^2 Y_{\mathbf{k}}, \quad \kappa(\mathbf{k}) = \sqrt{k(k+2)}.$$

On spatially compact FRW spacetimes, all homogeneous pure quasifree states can be represented in a particular form, and this will be the starting point for the definition of SLEs. To this end, we quote the following result.

3.1 Theorem. (Lüders and Roberts [15], Olbermann [16])

(A) Suppose that ω is a homogeneous pure quasifree state on $\mathcal{F}(M, g)$ for a spatially compact FRW spacetime with scale factor $a(t)$ and Hubble function $H(t)$.

Then there exists a sequence $\{T_k\}$, $k \in \mathbb{N}_0$, of C^∞ functions $T_k : \mathbb{R} \rightarrow \mathbb{C}$ which are solutions of the differential equations (where a dot indicates differentiation with respect to t and overlining means complex conjugation)

$$\ddot{T}_k(t) + 3H(t)\dot{T}_k(t) + \omega_k^2(t)T_k(t) = 0 \quad (t \in \mathbb{R}) \quad (4)$$

with the time-dependent frequencies

$$\omega_k^2(t) = \kappa(k)/a^2(t) + m^2 \quad (5)$$

and which fulfill the condition

$$\overline{T}_k(t)\dot{T}_k(t) - \dot{\overline{T}}_k(t)T_k(t) = ia^3(t) \quad (t \in \mathbb{R}), \quad (6)$$

such that, in the sense of distributions, one has

$$\mathcal{W}_2^\omega((t, \mathbf{x}), (t', \mathbf{x}')) = \sum_{\mathbf{k}=(k,l,m)} \bar{T}_k(t) Y_{\mathbf{k}}(\mathbf{x}) T_k(t') \bar{Y}_{\mathbf{k}}(\mathbf{x}'). \quad (7)$$

Relation (7) can equivalently be expressed by stating that the one-particle real-linear map $K_\omega : C_0^\infty(M, \mathbb{R}) \rightarrow \mathcal{H}_\omega^{(1)}$ is given as follows:

$\mathcal{H}_\omega^{(1)} = \ell^2 =$ space of square-summable sequences $\{s_{\mathbf{k}}\}$ indexed by the $\mathbf{k} = (k, l, m)$,

$K_\omega(F) = \{K_\omega(F)_{\mathbf{k}}\}$ with

$$K_\omega(F)_{\mathbf{k}} = \int_{\mathbb{R}} \int_{\Sigma} T_k(t) \bar{Y}_{\mathbf{k}}(\mathbf{x}) F(t, \mathbf{x}) dt d\mu_h(\mathbf{x}). \quad (8)$$

(B) Conversely, let $\{T_k\}$, $k \in \mathbb{N}_0$ be a sequence of smooth functions $T_k : \mathbb{R} \rightarrow \mathbb{C}$ fulfilling (4) and (6). Then the right hand side of (7) defines the two-point function $\mathcal{W}_2^\omega$ of a homogeneous pure quasifree state ω on $\mathcal{F}(M, g)$ for a spatially compact FRW spacetime with scale factor $a(t)$.

Thus, there is a one-to-one correspondence between the set of homogeneous pure quasifree states ω and the set of sequences $\{T_k\}$ of smooth functions fulfilling (4) and (6).²

Let us, from now on, denote this set of sequences by $\mathcal{T}$. We shall next quote a result by Olbermann which ultimately introduces states of low energy. These make reference to weighting functions. We shall say that a function $f : \mathbb{R} \rightarrow \mathbb{R}$ is a *weighting function* if there is a function $q \in C_0^\infty(\mathbb{R}, \mathbb{R})$ such that $f(t) = q(t)^2$ ($t \in \mathbb{R}$).

3.2 Theorem. (Olbermann [16])

For any weighting function f and any $\{T_k\} \in \mathcal{T}$, define the sequence of numbers

$$\tau_j(f, \{T_k\}) = \int_{\mathbb{R}} f(t) (|\dot{T}_j(t)|^2 + \omega_j^2(t) |T_j(t)|^2) dt \quad (j \in \mathbb{N}_0). \quad (9)$$

(i) Suppose that, for fixed weighting function f , the sequence $\{\overset{\circ}{T}_k\} \in \mathcal{T}$ has the minimizing property

$$\tau_j(f, \{\overset{\circ}{T}_k\}) = \min_{\{T_k\} \in \mathcal{T}} \tau_j(f, \{T_k\}) \quad (10)$$

for all $j \in \mathbb{N}_0$. Then the corresponding homogeneous pure quasifree state $\overset{\circ}{\omega} = \overset{\circ}{\omega}_f$ on $\mathcal{F}(M, g)$ is a Hadamard state. Furthermore, this homogeneous pure quasifree Hadamard state minimizes the energy density weighted by f along cosmological time, i.e.

$$\varrho_{\overset{\circ}{\omega}}[f] = \min_{\omega} \varrho_{\omega}[f] \quad (11)$$

where the minimum is taken over all homogeneous pure quasifree Hadamard states ω , with

$$\varrho_{\omega}[f] = \int_{\mathbb{R}} f(t) \langle \varrho : (t, \mathbf{x}) \rangle_{\omega} dt \quad (12)$$

²States which here are called “homogeneous” are called “isotropic” in [16].

for any $\mathbf{x} \in \Sigma$. (Owing to homogeneity of ω , the quantity on the right hand side of (12) is independent of $\mathbf{x}$.)

- (ii) For given weighting function f , a minimizing sequence $\{\overset{\circ}{T}_k\} \in \mathcal{T}$ exists. This sequence is unique up to choice of a phase factor for each k , i.e. $\{\overset{\circ}{T}_k\}$ has the minimizing property exactly if this is the case for $\{e^{ip_k}\overset{\circ}{T}_k\}$ for any sequence of real numbers $\{p_k\}$. All choices of phases lead to the same minimizing state $\overset{\circ}{\omega}$.

Given any sequence $\{S_k\} \in \mathcal{T}$, a minimizing sequence $\{\overset{\circ}{T}_k\}$ can be constructed from $\{S_k\}$ in the following way: One sets

$$\overset{\circ}{T}_k(t) = \lambda(k)S_k(t) + \mu(k)\overline{S}_k(t), \quad (13)$$

with the definitions

$$\begin{aligned} \mu(k) &= \sqrt{\frac{c_1(k)}{2\sqrt{c_1(k)^2 - |c_2(k)|^2}}} - \frac{1}{2} \\ \lambda(k) &= e^{i\xi_k} \sqrt{\frac{c_1(k)}{2\sqrt{c_1(k)^2 - |c_2(k)|^2}}} + \frac{1}{2} \\ c_1(k) &= \frac{1}{2} \int f(t)(|\dot{S}_k(t)|^2 + \omega_k^2(t)|S_k(t)|^2) dt \\ c_2(k) &= \frac{1}{2} \int f(t)(\dot{S}_k(t)^2 + \omega_k^2(t)S_k(t)^2) dt \\ \xi_k &= \pi - \text{Arg } c_2(k) \end{aligned} \quad (14)$$

Remarks

(α) Note that $\{\overset{\circ}{T}_k\}$, respectively $\overset{\circ}{\omega}$, minimize $\varrho_\omega[f]$ no matter how the renormalization constants A, B, Γ, D have been chosen.

(β) For fixed $\mathbf{x} \in \Sigma$, $\int_{\mathbb{R}} f(t) \langle : \varrho : (t, \mathbf{x}) \rangle_\omega dt$ equals the f -weighted integral of the expected energy density along the geodesic $t \mapsto (t, \mathbf{x})$ in FRW spacetime. It is known that this weighted integral is bounded below as a functional on quasifree Hadamard states ω . This fact is a special case of a quantum energy inequality, which has been established for the minimally coupled scalar field by Fewster [6].

(γ) Note also that $\lambda(k)$ and $\mu(k)$ and hence the minimizing sequence $\{\overset{\circ}{T}_k\}$ are unchanged under a constant rescaling $f(t) \mapsto r \cdot f(t)$ ($r > 0$) of the amplitude of the weighting function.

4 Bogoliubov transformations and particle creation

We continue to consider the case of spatially compact FRW spacetime (M, g) with scale factor $a(t)$ and Hubble function $H(t)$.

For any pure homogeneous quasifree state ω on $\mathcal{F}(M, g)$, the one-particle map $K_\omega : C_0^\infty(M, \mathbb{R}) \rightarrow \mathcal{H}_\omega^{(1)}$ has $\mathcal{H}_\omega^{(1)} = \ell^2$, with $K_\omega(\mathbf{f})$ given by (8); ℓ^2 is

identified as the space of square summable sequences $\{s_{\mathbf{k}}\}$, $\mathbf{k} = (k, l, m)$. The field operators $\Phi_{\omega}(\mathbf{F}) = \pi_{\omega}(\phi(\mathbf{F}))$ in the GNS-representation of ω take the form

$$\Phi_{\omega}(\mathbf{F}) = a(K_{\omega}(\mathbf{F})) + a^+(K_{\omega}(\mathbf{F})) \quad (15)$$

by (2) and (3). Let us denote by $\{\delta_{\mathbf{k}', \mathbf{k}}\}$ the sequence in ℓ^2 which takes the value 1 exactly if $\mathbf{k} = \mathbf{k}'$, and the value 0 else. Then write $a_{\mathbf{k}'} = a(\delta_{\mathbf{k}', \mathbf{k}})$ and $a_{\mathbf{k}'}^+ = a^+(\delta_{\mathbf{k}', \mathbf{k}})$. With this notation, and writing more simply $\mathbf{k}$ instead of $\mathbf{k}'$, one may recast (15) in the form

$$\Phi_{\omega}(t, \mathbf{x}) = \sum_{\mathbf{k}=(k,l,m)} (T_k(t) \bar{Y}_{\mathbf{k}}(\mathbf{x}) a_{\mathbf{k}}^+ + \bar{T}_k(t) Y_{\mathbf{k}}(\mathbf{x}) a_{\mathbf{k}}), \quad (16)$$

to be interpreted as an operator-valued distribution (the right hand side becomes an operator in Fock space $F_+(\mathcal{H}_{\omega}^{(1)})$ upon integrating with a test-function $\mathbf{F}(t, \mathbf{x})$).

It is known that for any pair of pure quasifree Hadamard states $\omega(1)$ and $\omega(2)$ on spatially compact FRW spacetime there is a unitary operator $U : F_+(\mathcal{H}_{\omega(1)}^{(1)}) \rightarrow F_+(\mathcal{H}_{\omega(2)}^{(1)})$ such that

$$\Phi_{\omega(2)}(\mathbf{F}) = U \Phi_{\omega(1)}(\mathbf{F}) U^{-1} \quad (17)$$

for all $\mathbf{F} \in C_0^\infty(M, \mathbb{R})$ [25, 22].

Now let $\omega(1)$ and $\omega(2)$ be two pure homogeneous quasifree Hadamard states and denote by $\{T_k^{(1)}\}$ and $\{T_k^{(2)}\}$ the corresponding sequences in $\mathcal{T}$. Since $T_k^{(1)}$ and $T_k^{(2)}$ both fulfill the differential equation (4) and the solution space of (4) is two-dimensional, there are complex coefficients α_k and β_k so that

$$T_k^{(2)}(t) = \alpha_k T_k^{(1)}(t) + \beta_k \bar{T}_k^{(1)}(t) \quad (t \in \mathbb{R}).$$

The requirement (6) implies

$$|\alpha_k|^2 - |\beta_k|^2 = 1.$$

Insertion in to (16) then yields the relations

$$\begin{aligned} \Phi_{\omega(1)}(t, \mathbf{x}) &= \sum_{\mathbf{k}=(k,l,m)} (T_k^{(1)}(t) \bar{Y}_{\mathbf{k}}(\mathbf{x}) a_{\mathbf{k}}^+ + \bar{T}_k^{(1)}(t) Y_{\mathbf{k}}(\mathbf{x}) a_{\mathbf{k}}) \quad \text{and} \\ \Phi_{\omega(2)}(t, \mathbf{x}) &= \sum_{\mathbf{k}=(k,l,m)} (T_k^{(2)}(t) \bar{Y}_{\mathbf{k}}(\mathbf{x}) b_{\mathbf{k}}^+ + \bar{T}_k^{(2)}(t) Y_{\mathbf{k}}(\mathbf{x}) b_{\mathbf{k}}) \end{aligned}$$

with

$$b_{\mathbf{k}}^+ = \alpha_k a_{\mathbf{k}}^+ + \bar{\beta}_{\mathbf{k}} a_{-\mathbf{k}}, \quad (18)$$

$$b_{\mathbf{k}} = \bar{\alpha}_{\mathbf{k}} a_{\mathbf{k}} + \beta_k a_{-\mathbf{k}}^+, \quad (19)$$

where the notation $-\mathbf{k} = (k, l, -m)$ for $\mathbf{k} = (k, l, m)$ and the property $\bar{Y}_{\mathbf{k}} = Y_{-\mathbf{k}}$ have been used. The passage from annihilation and creation operators $a_{\mathbf{k}}$ and $a_{\mathbf{k}}^+$ to $b_{\mathbf{k}}$ and $b_{\mathbf{k}}^+$ by (18), (19) is a special case of a *Bogoliubov transformation*. It is related to the unitary U appearing in (17) as formulated in the following,

well-known theorem [20, 8, 26, 18]:

There is a unitary $U : F_+(\mathcal{H}_{\omega(1)}^{(1)}) \rightarrow F_+(\mathcal{H}_{\omega(2)}^{(1)})$ such that

$$\Phi_{\omega(2)}(F) = U \Phi_{\omega(1)}(F) U^{-1} \quad (F \in C_0^\infty(M, \mathbb{R}))$$

exactly if

$$b_{\mathbf{k}} = U a_{\mathbf{k}} U^{-1} \quad \text{and} \quad b_{\mathbf{k}}^+ = U a_{\mathbf{k}}^+ U^{-1}$$

for all $\mathbf{k}$. This, in turn, holds if and only if

$$\sum_{\mathbf{k}=(k,l,m)} |\beta_k|^2 = \sum_{k \in \mathbb{N}_0} (k^2 + 2k + 1) |\beta_k|^2 < \infty. \quad (20)$$

Now suppose that we consider the GNS-representation of $\omega(2)$ as “reference” representation. With respect to this representation, we have $b_{\mathbf{k}}|0\rangle_{\omega(2)} = 0$, so the $b_{\mathbf{k}}$ and $b_{\mathbf{k}}^+$ are annihilation and creation operators of the spatial symmetry mode $\mathbf{k}$ with respect to this representation. One refers to this property also by saying that $b_{\mathbf{k}}$ and $b_{\mathbf{k}}^+$ annihilate and create “b-particles”. Correspondingly, the number operator

$$n_{\mathbf{k}}^{\omega(2)} = b_{\mathbf{k}}^+ b_{\mathbf{k}}$$

counts the number of of $\mathbf{k}$ -modes with respect to having chosen the GNS-representation of $\omega(2)$ as reference representation. One abbreviates that by saying that $n_{\mathbf{k}}^{\omega(2)}$ counts the number of $\mathbf{k}$ -modes of b-particles. Thus, $n_{\mathbf{k}}^{\omega(2)}$ can be regarded as a counter for $\mathbf{k}$ -modes which has been calibrated to give zero response in the reference state $\langle \cdot \rangle_{\omega(2)}$:

$$\langle n_{\mathbf{k}}^{\omega(2)} \rangle_{\omega(2)} = 0.$$

Then $n_{\mathbf{k}}^{\omega(2)}$ gives $|\beta_k|^2$ as response in the state $\langle \cdot \rangle_{\omega(2)}$:

$$\begin{aligned} \langle n_{\mathbf{k}}^{\omega(2)} \rangle_{\omega(1)} &= \omega(1) \langle 0 | b_{\mathbf{k}}^+ b_{\mathbf{k}} | 0 \rangle_{\omega(1)} \\ &= \omega(1) \langle 0 | (\alpha_k a_{\mathbf{k}}^+ + \bar{\beta}_k a_{-\mathbf{k}}) (\bar{\alpha}_k a_{\mathbf{k}} + \beta_k a_{-\mathbf{k}}^+) | 0 \rangle_{\omega(1)} \\ &= |\beta_k|^2 \end{aligned}$$

observing that $a_{\mathbf{k}}|0\rangle_{\omega(1)} = 0$ and $\omega(1) \langle 0 | a_{-\mathbf{k}}^+ a_{-\mathbf{k}} | 0 \rangle_{\omega(1)} = 1$.

We remark that in a situation where the scale factor $a(t)$ is never constant, it is impossible to have $|\beta_k|^2 = 0$ for all k unless the weighting functions f_1 and f_2 of the two SLEs $\omega(1)$ and $\omega(2)$ are proportional ($f_1(t) = r \cdot f_2(t)$ with a positive constant r). To see this, note that $\beta_k = 0$ implies that $b_{\mathbf{k}}|0\rangle_{\omega(1)} = 0$ which, in turn, means that $a_{\mathbf{k}} U^{-1} |0\rangle_{\omega(1)} = 0$. But the only non-zero vectors in $F_+(\mathcal{H}_{\omega(1)}^{(1)})$ which are annihilated by all $a_{\mathbf{k}}$ are scalar multiples of $|0\rangle_{\omega(1)}$. Likewise, the only non-zero vectors in $F_+(\mathcal{H}_{\omega(2)}^{(1)})$ which are annihilated by all the $b_{\mathbf{k}}$ are scalar multiples of $|0\rangle_{\omega(2)}$. Hence, $U|0\rangle_{\omega(1)} = e^{ir} |0\rangle_{\omega(2)}$ for some real r . By the properties of the GNS-representation, this implies that $\omega(1) = \omega(2)$ as states on $\mathcal{F}(M, g)$. However, Olbermann [16] has shown that for this to hold with non-constant $a(t)$, it is necessary that the weighting functions f_1 and f_2 are proportional. This feature for non-constant $a(t)$ is significantly different from

the case of constant $a(t) = a_0$, where there is one unique SLE for all weighting functions and hence always $\beta_k = 0$.

In the case of non-constant $a(t)$, the deviation of β_k from 0 is interpreted as (cosmological) particle creation. One may envisage the following situation portrayed in Figure 1: Some “initial” state $\omega(1)$ is prepared as SLE with respect to a weighting function f_1 concentrated near an early cosmological time t_1 . Another SLE $\omega(2)$ corresponding to a weighting function f_2 concentrated at later cosmological time $t_2 > t_1$ is used as reference state. This reference state can be taken as an approximate vacuum for an observer making measurements near the time t_2 . If the observer uses a particle (mode) counter calibrated to give zero response in his “approximate vacuum” $\omega(2)$, he or she will find that the symmetry mode $\mathbf{k}$ measured on the state $\omega(1)$ is excited with a distribution

$$\langle \mathbf{n}_{\mathbf{k}}^{\omega(2)} \rangle_{\omega(1)} = |\beta_k|^2.$$

On the other hand, another observer making measurements around the time t_1 using $\omega(1)$ as reference state will find no particle excitations when making measurements on the state $\omega(1)$. The way in which this “particle creation” depends on reference states that are defined with respect to properties localized in time should be noticed here.




Figure 1. *Illustration of the localizations of the weighting functions f_1 and f_2 in an exponentially expanding Universe. The parameters δ and ϵ denote the characteristic widths of these functions. The curve γ depicts the timelike geodesic of an inertial observer.*

5 Calculation of the number of created particles

As described in the previous section, we want to calculate the number of created particles per mode $\mathbf{k}$. We will make some special assumptions with regard to the Hubble function $H(t)$ and the mass parameter m and some further approximations for the sole purpose of facilitating the numerical calculations. Let us explain how we proceed in several steps.

(I)

Supposing that a pair of weighting functions f_1 and f_2 has been chosen, we denote by $\omega(1)$ and $\omega(2)$ the corresponding SLEs (pure, homogeneous, quasifree) and by $\{T_k^{(1)}\}$ and $\{T_k^{(2)}\}$ the corresponding sequences in $\mathcal{S}$. They are related by the Bogoliubov coefficients α_k and β_k ($k \in \mathbb{N}_0$) via

$$T_k^{(2)}(t) = \alpha_k T_k^{(1)}(t) + \beta_k \overline{T}_k^{(1)}(t) \quad (t \in \mathbb{R}, k \in \mathbb{N}_0) \quad (21)$$

Using moreover (cf. (6))

$$\overline{T}_k^{(j)}(t) \dot{T}_k^{(j)}(t) - \dot{\overline{T}}_k^{(j)}(t) T_k^{(j)}(t) = ia(t)^{-3} \quad (t \in \mathbb{R}, k \in \mathbb{N}_0) \quad (22)$$

where $a(t)$ is the scale factor of the underlying FRW spacetime, one finds that (21) and (22) imply

$$\beta_k = ia^3 \left(\dot{T}_k^{(2)} T_k^{(1)} - \dot{T}_k^{(1)} T_k^{(2)} \right) \Big|_{t^*} \quad (23)$$

for any choice of $t^* \in \mathbb{R}$.

(II)

The best way to find $\{T_k^{(1)}\}$ and $\{T_k^{(2)}\}$ is to choose some $\{S_k\} \in \mathcal{T}$ and to calculate the coefficients $\lambda^{(j)}(k)$ and $\mu^{(j)}(k)$ ($j = 1, 2$) as in (...) so that

$$T_k^{(j)}(t) = \lambda^{(j)}(k) S_k(t) + \mu^{(j)}(k) \overline{S}_k(t). \quad (24)$$

Observing that

$$\overline{S}_k \dot{S}_k(t) - \dot{\overline{S}}_k(t) S_k(t) = ia(t)^{-3}, \quad (25)$$

equations (23), (24) and (25) combine to yield

$$\beta_k = \lambda^{(1)}(k) \mu^{(2)}(k) - \lambda^{(2)}(k) \mu^{(1)}(k) \quad (26)$$

for all $k \in \mathbb{N}_0$.

(III)

We will now specify the Hubble function as $H(t) = H$ with a constant $H > 0$, thus

$$a(t) = e^{Ht}, \quad (27)$$

and we will choose the unit of time such that $H = 1$. In view of the fact that we have assumed our underlying FRW spacetime to be spatially closed, this scenario doesn't comply with a solution of the Einstein equations with normal matter (ideal fluid) without cosmological constant; rather it models an epoch of accelerated expansion of the Universe that may have taken place over a time-span such that $H = 1$. For the purposes of illustration, we will nevertheless take

that time-span here to be the estimated age of the Universe, $t_H = 1.39 \cdot 10^{10}y$. (One may take any other (shorter) time-span upon scaling the mass parameter m accordingly to interpret our numerical results analogously, see below.) Writing as before $\kappa = \sqrt{k(k+2)}$ and inserting (27), the S_k must satisfy (4), which now takes the form

$$\ddot{S}_k(t) + 3H\dot{S}_k(t) + (\kappa^2 e^{-2Ht} + m^2)S_k(t) = 0. \quad (28)$$

With the unit of time chosen so that $H = 1$, these equations simplify further as

$$\ddot{S}_k(t) + 3\dot{S}_k(t) + (\kappa^2 e^{-2t} + m^2)S_k(t) = 0. \quad (29)$$

(IV)

For each $\kappa = \sqrt{k(k+2)}$, the differential equation (29) has two linearly independent solutions, given by

$$y_{\kappa}^{\pm\nu}(t) = e^{-\frac{3}{2}t} J_{\pm\nu}(e^{-t}\kappa), \quad (30)$$

where the order of the Bessel functions is

$$\nu = \frac{\sqrt{9 - 4m_*^2}}{2} \quad (31)$$

with m_* the numerical value of the mass parameter m in the units employed, i.e. $m_* = c^2 m / (\hbar H)$ when m is given in cgs units. The Bessel functions specialize in case that $\nu = \pm 1/2$ according to

$$J_{\frac{1}{2}}(x) = \sqrt{\frac{2}{\pi x}} \sin x \quad J_{-\frac{1}{2}}(x) = \sqrt{\frac{2}{\pi x}} \cos x \quad (32)$$

for real x , and since this situation simplifies the numerical calculation of particle creation considerably, we will assume that m has been chosen such that $\nu = \pm 1/2$. This corresponds to $m_* = \sqrt{2}$, or $m = \sqrt{2}\hbar H / c^2 \approx 3 \times 10^{-69} \text{kg}$ in cgs units. This is surely a very small mass, but again, this value is chosen for the purpose of illustration so as to make the numerical calculations easier. With that choice of m and correspondingly $m_* = \sqrt{2}$, and the identities (32), one obtains for each $k \in \mathbb{N}_0$ a solution

$$S_k(t) = A(\kappa)e^{-t} \cos(e^{-t}\kappa) + B(\kappa)e^{-t} \sin(e^{-t}\kappa) \quad (33)$$

to (29), where the coefficients

$$A(\kappa) = \frac{(2i+1)\sin\kappa}{2\kappa} + \frac{\cos\kappa}{2} \quad \text{and} \quad (34)$$

$$B(\kappa) = \frac{-(2i+1)\cos\kappa}{2\kappa} + \frac{\sin\kappa}{2}. \quad (35)$$

have been chosen such that the initial conditions

$$\dot{S}_k(0) = i, \quad S_k(0) = \frac{1}{2}, \quad (36)$$

are fulfilled. This ensures that the $S_k(t)$ defined in (33) fulfill the normalization condition (24), and therefore give rise to a sequence $\{S_k\} \in \mathcal{T}$.

(V)

Now that we have chosen some $\{S_k\} \in \mathcal{T}$, we will calculate the Bogoliubov coefficients as described in **(II)** above. To this end, we must evaluate the integrals

$$c_1^{(i)}(\kappa) = \frac{1}{2} \int dt f_i(t) \left(|\dot{S}_k(t)|^2 + \omega_k^2 |S_k(t)|^2 \right)$$

and

$$c_2^{(i)}(\kappa) = \frac{1}{2} \int dt f_i(t) \left(\dot{S}_k(t)^2 + \omega_k^2 S_k(t)^2 \right) ,$$

which depend on the smearing functions $f_i(t)$ ($i = 1, 2$) and $S_k(t)$. After some algebra one can write the integrands as

$$\begin{aligned} |\dot{S}_k(t)|^2 + \omega_k^2 |S_k(t)|^2 = & \\ & \left(\frac{5}{4} + \frac{\kappa^2}{4} \right) I_1 + \left(\frac{3}{8} + \frac{15}{8\kappa^2} \right) I_2 - \frac{1}{2} I_3 + \left(\frac{3}{8} - \frac{15}{8\kappa^2} \right) I_4 \\ & + \left(\frac{\kappa}{4} - \frac{5}{4\kappa} \right) I_5 + \frac{3}{4\kappa} I_6 \end{aligned} \quad (37)$$

and

$$\begin{aligned} \dot{S}_k(t)^2 + \omega_k^2 S_k(t)^2 = & \\ & \left(i - \frac{3}{4} + \frac{\kappa^2}{4} \right) I_1 + \left(\frac{3}{8} - \frac{9}{8\kappa^2} + \frac{3i}{2\kappa^2} \right) I_2 - \left(i + \frac{1}{2} \right) I_3 \\ & + \left(\frac{3}{8} + \frac{9}{8\kappa^2} - \frac{3i}{2\kappa^2} \right) I_4 + \left(\frac{3}{4\kappa} - \frac{i}{\kappa} + \frac{\kappa}{4} \right) I_5 + \left(\frac{3}{4\kappa} + \frac{3i}{2\kappa} \right) I_6 , \end{aligned} \quad (38)$$

where the notation

$$\begin{aligned} I_1 &= e^{-4t} & I_4 &= e^{-2t} \cos \left[(2 - 2e^{-t}) \kappa \right] \\ I_2 &= e^{-2t} & I_5 &= e^{-3t} \sin \left[(2 - 2e^{-t}) \kappa \right] \\ I_3 &= e^{-3t} \cos \left[(2 - 2e^{-t}) \kappa \right] & I_6 &= e^{-2t} \sin \left[(2 - 2e^{-t}) \kappa \right] \end{aligned}$$

has been used.

(VI)

As weighting functions $f_i(t)$ we would have to use squares of C_0^∞ -functions. However, since we are eventually evaluating the integrals $c_1^{(i)}$ and $c_2^{(i)}$ numerically on a computer, it is justified to approximate (up to machine precision) weighting functions by Gaussians. Therefore, for our numerical calculations we choose as weighting functions $f_i(t)$ Gaussians localised at the times t_i and characteristic smearing widths ϵ and δ as illustrated in Figure 2 (cf. also Figure 1):

$$f_1(t) = \frac{1}{\epsilon} e^{-(\frac{t-t_1}{\epsilon})^2} , \quad f_2(t) = \frac{1}{\delta} e^{-(\frac{t-t_2}{\delta})^2} . \quad (39)$$




Figure 2. *Illustration of the parameters t_1, t_2, ϵ and δ which characterize the weighting functions f_1 and f_2 .*

As said, this is a sufficiently good approximation for a weighting function as long as the smearing widths ϵ and δ are much smaller than the unit of time (i.e. much smaller than 1 with respect to our choice of units). The normalisation factors $1/\epsilon$ and $1/\delta$ have no impact on the $\mu^{(i)}(\kappa)$ and $\lambda^{(i)}(\kappa)$ since they will cancel. Nevertheless we wrote them down for one can better imagine the limit to the Dirac distribution when ϵ and δ tend towards zero. The time coordinate is not subject to any restrictions a priori. Therefore the integrals

$$\mathcal{I}_j(\epsilon, t_1) = \int_{-\infty}^{\infty} dt f_1(t) I_j \quad (j = 1, \dots, 6)$$

have to be evaluated. In order to apply the identities

$$\int_{-\infty}^{\infty} dx e^{-q^2 x^2} \sin[p(x+y)] = \frac{\sqrt{\pi}}{q} e^{-\frac{p^2}{4q^2}} \sin[py] \quad (40)$$

$$\int_{-\infty}^{\infty} dx e^{-q^2 x^2} \cos[p(x+y)] = \frac{\sqrt{\pi}}{q} e^{-\frac{p^2}{4q^2}} \cos[py] \quad (41)$$

for the integrations, we carry out the Taylor expansion of the argument of the trigonometric functions to first order in $t - t_1$. This approximation is justified if $\epsilon \ll 1$, since then the integrand will only be important for $|t - t_1| \ll 1$. We obtain

$$2\kappa(1 - e^{-t}) \approx 2\kappa(1 - e^{-t_1}) + \frac{2\kappa}{e^{t_1}}(t - t_1). \quad (42)$$

Multiplication of the two exponential functions yields

$$e^{-(\frac{t-t_1}{\epsilon})^2} e^{-\alpha t} = e^{-\frac{1}{\epsilon^2}(t-(t_1-\frac{\alpha}{2}\epsilon^2))^2} e^{-\alpha t_1} \quad (43)$$

where we used $(\alpha\epsilon)^2 \ll 1$. We define a new integration variable $\tau := t - (t_1 - \frac{\alpha}{2}\epsilon^2)$ and rewrite the argument of the trigonometric function:

$$2\kappa(1 - e^{-Ht_1}) + \frac{2\kappa}{e^{t_1}}(t - t_1) = \frac{2\kappa}{e^{t_1}}\left(\tau - \frac{\alpha\epsilon^2}{2} - 1 + e^{t_1}\right) \quad (44)$$

In our case we have

$$\begin{aligned} q &= \frac{1}{\epsilon} , & y &= -\frac{\alpha}{2}\epsilon^2 + e^{t_1} - 1 \\ p &= \frac{2\kappa}{e^{t_1}} , & py &\approx 2\kappa(1 - e^{-t_1}) . \end{aligned} \quad (45)$$

Consequently,

$$\begin{aligned} \mathcal{J}_1(\epsilon, t_1) &= \sqrt{\pi} e^{-4t_1} \\ \mathcal{J}_2(\epsilon, t_1) &= \sqrt{\pi} e^{-2t_1} \\ \mathcal{J}_3(\epsilon, t_1) &= \sqrt{\pi} e^{-3t_1} e^{-\frac{\kappa^2 \epsilon^2}{\exp(2t_1)}} \cos[2\kappa(1 - e^{-t_1})] \\ \mathcal{J}_4(\epsilon, t_1) &= \sqrt{\pi} e^{-2t_1} e^{-\frac{\kappa^2 \epsilon^2}{\exp(2t_1)}} \cos[2\kappa(1 - e^{-t_1})] \\ \mathcal{J}_5(\epsilon, t_1) &= \sqrt{\pi} e^{-3t_1} e^{-\frac{\kappa^2 \epsilon^2}{\exp(2t_1)}} \sin[2\kappa(1 - e^{-t_1})] \\ \mathcal{J}_6(\epsilon, t_1) &= \sqrt{\pi} e^{-2t_1} e^{-\frac{\kappa^2 \epsilon^2}{\exp(2t_1)}} \sin[2\kappa(1 - e^{-t_1})] , \end{aligned} \quad (46)$$

which completes our task of calculating the integrals $c_1^{(i)}$ and $c_2^{(i)}$ for $i = 1$. For $i = 2$ the same procedure as described above applies. We just have to perform the substitutions $t_1 \rightarrow t_2$ and $\epsilon \rightarrow \delta$ and to carry over the corresponding assumption $\delta \ll 1$.

(VII)

The explicit calculation of $|\beta_k|^2$ according to equation (26) is elementary but very cumbersome. Below, we contend ourselves with calculation of $|\beta_k|^2$ for several constellations of the parameters ϵ, t_1, δ and t_2 . The diagrams were generated with the program “Mathematica”. The particle creation coefficients $|\beta(\kappa)|^2 = |\beta_k|^2$ are plotted against $\kappa = \sqrt{k(k+2)}$ continuously, although actually κ is discrete in our case where $\Sigma = S^3$. Furthermore, all times are represented in the natural unit $t_H = 1, 3787 \times 10^{10} \text{y}$.

We briefly recall our assumptions made for our numerical calculations:

- The scale factor is $a(t) = e^t$.
- The mass of the particle has numerical value $m_* = \sqrt{2}$ in the units used.
- The test functions are approximated by Gaussians localised around t_1, t_2 and have characteristic smearing widths ϵ, δ .
- $\epsilon, \delta \ll 1$

5.1 The case $t_1 = t_2$

We consider the case $t_1 = t_2$ and investigate the influence of ϵ and δ on the particle creation. The special case $\epsilon = \delta$ is trivial: There is no particle creation since $\omega(1) = \omega(2)$




Figure 3. Particle creation rate for $t_1 = t_2 = 0.1$, $\delta = 10^{-5}$ for various values of ϵ with $\epsilon \approx \delta$

As soon as ϵ and δ deviate minimally ($\epsilon \approx \delta$), small variations of ϵ at constant δ will affect the particle creation rate $|\beta(\kappa)|^2$ in the region $\kappa = e^{t_1}\pi/(4\epsilon) \approx e^{t_1}\pi/(4\delta)$ (see Figure 3). Thus the magnitude of smearing widths determines which modes will be excited. As another case we consider a scale difference between ϵ and δ ($\epsilon \ll \delta$), cf. Figure 4. We remark that the maximum of the curve $|\beta(\kappa)|^2$ conforms to the larger smearing width (which is ϵ in our case) according to the formula

$$\kappa_{\max} \approx \frac{e^{t_1}\pi}{4\delta} \quad (47)$$

Small variations of ϵ influence $|\beta(\kappa)|^2$ only around $\kappa_{\max}$.




Figure 4. Particle creation rate for $t_1 = t_2 = 0.1$, $\delta = 10^{-5}$ and various values of ϵ for the case $\epsilon \gg \delta$




Figure 5. Particle creation rate for $\epsilon = 2\delta = 2 \times 10^{-5}$ and logarithmically increasing values of $t_1 = t_2$

Now we look at the effects caused by increasing values of $t_1 = t_2$, keeping ϵ and δ fixed (Figure 5). With logarithmically growing preparation times, $\kappa_{\max}$ increases linearly, while $|\beta(\kappa_{\max})|^2$ remains constant. Simultaneously the curve broadens. This is due to the properties of the solutions $S_k(t)$ in an exponentially increasing Universe where the frequency of oscillations decays $\propto e^{-t}$. Thus the ratio of ϵ and δ (which remain constant) to the period of the oscillations decreases, leading to the same effect as shifting $\kappa_{\max}$ to the right which would be obtained by decreasing ϵ and δ at constant t_1, t_2 , cf. (47). Physically this corresponds to the redshift of a particle associated with the mode κ which loses energy. Thus at later times the same mode will be excited more easily. In the diagram this is represented by the fact that curves corresponding to later times arise from those at earlier times by a κ -depending dilation to the right.

Finally we remark that the common feature of the case $t_1 = t_2$ is $|\beta(0)|^2 = 0$.

5.2 The case $t_1 \neq t_2$

We consider now the (more general) situation when t_1 and t_2 are different. Due to the difference $t_2 - t_1$ there are phase differences in the trigonometric functions entering in $|\beta(\kappa)|^2$, which manifest themselves in oscillations of the particle creation curve as seen in Figure 6. Their frequencies grow with increasing time difference $t_2 - t_1$. Due to these phase differences, $|\beta(0)|^2$ is no longer equal to zero. Thus, apart from the oscillation effects there is another new remarkable feature: the particle creation effect is biggest for small modes. When imposing much larger time differences $t_2 - t_1$, the particle creation curve grows and oscillates with higher frequency (Figure 7).




Figure 6. *Particle creation rate for $\epsilon = \delta = 10^{-5}$, $t_1 = 0.1$ and various values of t_2 corresponding to a small time difference*

There is some similarity with findings in [11], where it is also argued that the particle number of a state prepared at early times can diminish when compared with a “vacuum state” at later time. However, the authors of [11] use instantaneous vacuum states as reference states, which is problematic since they aren’t (locally) unitarily equivalent to Hadamard states [9, 12].




Figure 7. *Envelopes of the particle creation rate for $\epsilon = \delta = 10^{-5}$, $t_1 = 0.1$ and various values of t_2 corresponding to a big time difference. We have suppressed the oscillations because they are of too high frequency to be properly represented.*

We have already remarked towards the end of Sec. 4 that in the case $t_2 > t_1$, so that f_1 and f_2 aren’t proportional, it can’t happen that $|\beta(\kappa)|^2 = 0$ for all

values of κ . This is corroborated by the numerical results represented in Figures 6 and 7.

Another case of interest is the limit $\epsilon \rightarrow 0$ and/or $\delta \rightarrow 0$. In this case $f_1(t)$ and/or $f_2(t)$ would no longer be test functions, but delta distributions. Formally, the SLEs corresponding to the f_i would, in these limits, tend to instantaneous vacuum states, which fail to have the Hadamard property. This implies that these limiting states are no longer unitarily equivalent to any other SLE. This in turn should be expressed in the divergence of the total particle number in the corresponding squeezed vacuum vector, which is equivalent to $|\beta(\kappa)|^2 \geq O(\kappa^{-3})$. In fact one can show (see [4]) that in our example we have for fixed $\delta > 0$ in the limit $\epsilon = 0$,

$$|\beta(\kappa)|^2 \geq \frac{e^{2t_1}}{4\kappa^2} \sin^2(2(1 - e^{-t_1})\kappa) + O(\kappa^{-3}).$$

Comparing this with the criterion (20), one can see that the limiting state for $\epsilon \rightarrow 0$ isn't unitarily equivalent to the fixed SLE $\omega(2)$, and hence not a Hadamard state.

6 Summary and outlook

We have calculated the expected number $|\beta_k|^2$ of created particles per frequency mode k for an initial SLE at early cosmological time t_1 in a reference SLE at late cosmological time t_2 in a closed, exponentially expanding Universe. A characteristic feature is that $|\beta_k|^2$ shows oscillatory behaviour with respect to variation of k . The envelope of $|\beta_k|^2$ decays in k more strongly than $\sim k^{-3}$, so that the oscillations are most significant for low frequency modes. The oscillatory behaviour increases with growing time-difference $t_2 - t_1$.

A substantial drawback in discussing the possible physical significance of these findings is that they have been obtained under considerably simplifying assumptions which cannot really be considered as physically realistic. This refers mostly to having made the assumption of a spatially closed, exponentially expanding FRW spacetime and having fine-tuned the mass term m in the scalar field equation to an extremely small value so as to make the numerical calculation more tractable. Above that, the quantized linear scalar field is a toy model to which there corresponds no observed particle. It is, then, much desirable to extend the investigation of this article to other quantum field models such as the Dirac and electromagnetic fields, and for physically realistic cosmological scenarios and parameters.

However, the basic features of our methods should carry over also to these more realistic models, with qualitatively analogous results. Therefore, the main purpose of this paper is to show that cosmological particle creation can be rigorously discussed, and detailed numerical results can be obtained, when employing states of low energy as reference states. This, in turn, also demonstrates the utility of the class of SLEs in the context of cosmological considerations.

References

- [1] Bär, C., Ginoux, N, Pfäffle, F., *Wave Equations on Lorentzian Manifolds and Quantization*, European Mathematical Society, Zürich, 2007

- [2] Brunetti, R., Fredenhagen, K., Verch, R., “The Generally Covariant Locality Principle – A New Paradigm for Local Quantum Field Theory”, *Commun. Math. Phys.* **237** (2003) 31
- [3] Dappiaggi, C., Fredenhagen, K., Pinamonti, N., “Stable Cosmological Models Driven by a Free Quantum Scalar Field”, *Phys. Rev.* **D77** (2008) 104015
- [4] Degner, A., Diploma Thesis, Dept. of Physics, University of Leipzig, 2008
- [5] Dimock, J., “Algebras of Local Observables on a Manifold”, *Commun. Math. Phys.* **77** (1980) 219
- [6] Fewster, C.J., “A General Worldline Quantum Inequality”, *Class. Quantum Grav.* **17** (2000) 1897
- [7] Fewster, C.J., Smith, C.J., “Absolute Quantum Energy Inequalities in Curved Spacetime”, *Annales Henri Poincaré* **9** (2008) 425
- [8] Fulling, S.A., *Aspects of Quantum Field Theory in Curved Space-Time*, Cambridge University Press, Cambridge, 1989
- [9] Fulling, S.A., “Remarks on Positive Frequency and Hamiltonians in Expanding Universes”, *Gen. Rel. Grav.* **10** (1979) 807
- [10] Hollands, S., Wald, R.M., “Axiomatic Quantum Field Theory in Curved Spacetime”, e-Print arXiv:0803.2003 [gr-qc]
- [11] Hu, B.L., Kang, G., Matacz, A., “Squeezed Vacua and the Quantum Statistics of Cosmological Particle Creation”, *Int. J. Mod. Phys.* **A9** (1994) 991
- [12] Junker, W., “Hadamard States, Adiabatic Vacua and the Construction of Physical States for Scalar Quantum Fields on Curved Space-Time”, *Rev. Math. Phys.* **8** (1996) 1091; Erratum *ibid.* **14** (2002) 511
- [13] Junker, W., Schrohe, E., “Adiabatic Vacuum States on General Space-Time Manifolds”, *Annales Henri Poincaré* **3** (2002) 1113
- [14] Kay, B.S., Wald, R.M., “Theorems on the Uniqueness and Thermal Properties of Stationary, Nonsingular, Quasifree States on Space-Times with a Bifurcate Killing Horizon”, *Phys. Rep.* **207** (1991) 49
- [15] Lüders, C., Roberts, J.E., “Local Quasiequivalence and Adiabatic Vacuum States”, *Commun. Math. Phys.* **134** (1990) 29
- [16] Olbermann, H., “States of Low Energy on Robertson-Walker Spacetimes”, *Class. Quantum Grav.* **24** (2007) 5011
- [17] Parker, L., “Quantized Fields and Particle Creation in Expanding Universes. 1.”, *Phys. Rev.* **183** (1969) 1057
- [18] Ruijsenaars, S.N.M., “On Bogolyubov Transformations. 2. The General Case”, *Ann. Physics (N.Y.)* **116** (1978) 105
- [19] Schlemmer, J., Verch, R., “Local Thermal Equilibrium States and Quantum Energy Inequalities”, *Ann. Henri Poincaré* **9** (2008) 945

- [20] Shale, D., “Linear Symmetries of Free Boson Fields”, Trans. Amer. Math. Soc. **103** (1962) 149
- [21] Streater, R.F., Wightman, A.S., *PCT, Spin and Statistics, and All That*, Benjamin, New York, 1964
- [22] Verch, R., “Local Definiteness, Primarity and Quasiequivalence of Quasifree Hadamard Quantum States in Curved Spacetime”, Commun. Math. Phys. **160** (1994) 507
- [23] Wald, R.M., “The Back Reaction Effect in Particle Creation in Curved Spacetime”, Commun. Math. Phys. **54** (1977) 1
- [24] Wald, R.M., *General Relativity*, University of Chicago Press, Chicago, 1984
- [25] Wald, R.M., *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics*, University of Chicago Press, Chicago, 1994
- [26] Wald, R.M., “Existence of the S Matrix in Quantum Field Theory in Curved Spacetime”, Ann. Physics (N.Y.) **118** (1979) 490
- [27] Winitzki, S., “Cosmological Particle Production and the Precision of the WKB Approximation”, Phys. Rev. **D72** (2005) 104011